

19



Europäisches Patentamt
European Patent Office
Office européen des brevets

11 Publication number:

0 131 302
A2

12

EUROPEAN PATENT APPLICATION

21 Application number: **84108015.3**

51 Int. Cl.⁴: **C 07 D 233/60**
C 07 D 233/61, A 61 K 31/415

22 Date of filing: **09.07.84**

30 Priority: **12.07.83 US 513143**

43 Date of publication of application:
16.01.85 Bulletin 85/3

84 Designated Contracting States:
AT BE CH DE FR GB IT LI LU NL SE

71 Applicant: **SCHERING AKTIENGESELLSCHAFT Berlin und Bergkamen**
Müllerstrasse 170/178 Postfach 65 03 11
D-1000 Berlin 65(DE)

72 Inventor: **Diamond, Julius**
179 Lake Drive
Mountain Lakes 07046 New Jersey(US)

72 Inventor: **Lumma, William Carl, Jr.**
25 Newman Road, M.R. 1
Pennsburg 18073 Pennsylvania(US)

72 Inventor: **Morgan, Thomas Kenneth, Jr.**
47 Sherman Avenue
Morris Plains 07950 New Jersey(US)

72 Inventor: **Wohl, Ronald André**
41-2B Mt. Pleasant Village
Morris Plains 07950 New Jersey(US)

54 **Imidazoliums, intermediates thereto and their use.**

57 The imidazolium salts described are useful as antiarrhythmic agents. A method of treating arrhythmia by increasing the refractoriness of cardiac tissue is provided as well as pharmaceutical formulations containing such imidazolium salts.

EP 0 131 302 A2


IMIDAZOLIUMS, INTERMEDIATES
THERE TO AND THEIR USE


This invention relates to novel quaternary salts and their use as antiarrhythmic agents. Specifically, this invention relates to imidazolium salts and their molecular compounds with aromatic dicarboxylic acids, pharmaceutical compositions containing them as active ingredients and to the method of using them for the treatment of arrhythmia. This invention also relates to certain intermediate substituted imidazoles useful as antiarrhythmic agents and pharmaceutical compositions containing them as active ingredients.


GENERAL DESCRIPTION OF THE INVENTION

COMPOSITION-OF-MATTER ASPECT

In its composition-of-matter aspect, this invention relates to novel imidazolium salts. Particularly, this invention relates to the novel compounds defined by the following Formula I:



wherein R is a straight or branched chain alkyl having 1-12 carbon atoms, cycloalkyl(lower)alkyl, $-(\text{CH}_2)_t-$  R_8 , or



$\text{R}_1, \text{R}_2, \text{R}_3$ may be the same or independently, hydrogen, a straight chain alkyl group of 1-4 carbon atoms; R_4, R_5 may be the same or independently, hydrogen, halogen, hydroxy, lower alkyl, lower alkoxy, cyano, nitro, carbamoyl, lower alkylcarbamoyl, lower alkanoylamido, sulfamoyl, substituted sulfamoyl, sulfamoylamino.

N-lower alkylsulfamoylamino, lower alkylsulfonamido, substituted lower alkylsulfonamido, trifluoromethanesulfonamido, ureido, N-lower alkylureido, or when taken together on adjacent carbon atoms of the ring to which they are attached may be joined to form the methylenedioxy moiety;

R_6 is hydrogen, halogen, hydroxy, cyano, carbamoyl, lower alkoxy, straight chain alkyl of 1-6 carbon atoms;

A is phenyl, furanyl, pyrrolyl, pyridinyl, thiazoyl, oxazolyl, the benzo derivatives of the foregoing and thiadiazolyl;

n is an integer of 0-2;

p is an integer of 0-5;

t is an integer of 1-5;


Y is methylene, hydroxymethylene, arylmethylene, substituted arylmethylene, phenylene, ethynylene, $\begin{smallmatrix} R_7 & R_8 \\ | & | \\ -C & =C- \end{smallmatrix}$, $\begin{smallmatrix} R_9 & R_{10} \\ | & | \\ -CH & -CH- \end{smallmatrix}$,

$\begin{smallmatrix} R_7 & R_8 \\ \diagup & \diagdown \\ C \\ \diagdown & \diagup \end{smallmatrix}$, carbonyl, $>N-R_{11}$, $-NHSO_2-$, $-SO_2NH-$; wherein R_7 , R_8

is hydrogen, C_1-C_4 lower alkyl or taken together form a 5 or 6 membered ring; R_9 , R_{10} is C_1-C_4 lower alkyl, or taken together may form a 5 or 6 membered ring or either may be hydrogen if the other is C_1-C_4 alkyl; R_{11} is hydrogen, lower alkyl, cyano, carbamoyl; with the provisos that when:

(a) Y is $>N-R_{11}$, $-SO_2NH-$, $-NHSO_2-$, then p must be an integer of 2-5,

(b) Y is hydroxymethylene then R cannot be



(c) Y is methylene, ethynylene, $\begin{smallmatrix} R_7 & R_8 \\ | & | \\ -C & =C- \end{smallmatrix}$, $\begin{smallmatrix} & R_7 \\ & | \\ =C & =C \\ & | \\ & R_8 \end{smallmatrix}$

then p must be an integer of 1-5,

(d) the sum of the carbon atoms in the chain n, Y, p is 2
then both R₄ and R₅ cannot be hydrogen,

(e) Y is carbonyl then both R₄ and R₅ cannot be hydrogen;

Z⁻ = halide, tosylate, sulfate, phosphate, methanesulfonate,
and the molecular compounds of the foregoing quaternary salts
with aromatic dicarboxylic acids.

In the above Formula I, halogen represents fluorine, chlorine,
bromine and iodine. The terms lower alkyl, lower alkoxy are
taken to mean a straight or branched 1-4 carbon chain. Substituted
sulfamoyl shall refer to lower alkyl substitution on the sulfamoyl
nitrogen, which in turn may be substituted by a phenyl or lower
alkoxy group and substituted lower alkylsulfonamido shall be
taken to mean, phenyl or lower alkoxy substitution on the lower
alkyl moiety. N-lower alkyluriedo shall be taken to mean lower
alkyl—NH— $\overset{\text{O}}{\parallel}$ —NH—.

Preferred classes of compounds embodied by this invention
are those of the above general formula having one or more of the
following characteristics:

- (1) A is phenyl and one of R₄ or R₅ is other than hydrogen
- (2) R is C₇H₁₅
- (3) R is CH₃
- (4) R₃ is $\dot{\text{C}}\text{H}_3$

The following are some of the compounds which exemplify
various aspects of the invention described herein.

- (1) 1-[4-(4-Chlorophenyl)butyl]-3-heptylimidazolium chloride.
- (2) 1-[4-(4-Chlorophenyl)butyl]-3-heptylimidazolium dihydro-
genphosphate.
- (3) 1-(4-Methoxyphenylpropyl)-3-methylimidazolium bromide.
- (4) 1-Butyl-3-[3-(4-methoxyphenyl)propyl]imidazolium bromide.
- (5) 1-[4-(4-Chlorophenyl)butyl]-3-heptylimidazolium chloride
(1:1) compound with phthalic acid.
- (6) 1-Heptyl-3-[3-(4-methoxyphenyl)propyl]imidazolium bromide.

- (7) 1-Heptyl-3-[4-(methoxyphenyl)butyl]imidazolium bromide.
- (8) 1-Heptyl-3-[4-(4-methoxyphenyl)butyl]-2-methylimidazolium bromide.
- 5 (9) 1-[3-((2,6-Dimethylphenyl)sulfamoyl)propyl]-3-heptyl-imidazolium chloride.
- (10) (E)-1-[4-(4-Chlorophenyl)-2-butenyl]-3-heptylimidazolium chloride (1:1) compound with phthalic acid.
- (11) 1-[4-(2,3-Dichlorophenyl)butyl]-3-heptylimidazolium chloride.
- (12) 1-Heptyl-3-(4-phenylbutyl)imidazolium chloride 1:1 compound
10 with phthalic acid.
- (13) 1-[3-(4-Chlorophenyl)propyl]-3-heptylimidazolium chloride.
- (14) 1-[4-(4-(Acetylamino)phenyl)butyl]-3-heptylimidazolium 4-methylbenzenesulfonate.
- (15) 1-[4-(4-Chlorophenyl)butyl]-3-methylimidazolium chloride.
- 15 (16) 1-[4-(4-Chlorophenyl)butyl]-3-propylimidazolium dihydrogen-phosphate.
- (17) 1-[4,4-Diphenylbutyl]-3-heptylimidazolium dihydrogenphosphate.
- (18) 1-[4-(4-Chlorophenyl)butyl]-3-(cyclohexylmethyl)imidazolium dihydrogenphosphate.
- 20 (19) 1-[4-(4-Fluorophenyl)butyl]-3-heptylimidazolium dihydrogen-phosphate.
- (20) 1-[4-(4-Chlorophenyl)butyl]-3-pentylimidazolium dihydrogen-phosphate.
- (21) 1-Butyl-3-[4-(4-chlorophenyl)butyl]imidazolium dihydrogen-phosphate.
25
- (22) 1-Heptyl-3-[4-[4-[(methyl)sulfonyl]amino]phenyl]butyl]-imidazolium dihydrogenphosphate.
- (23) 1,3-bis [4-(4-Chlorophenyl)butyl]imidazolium dihydrogen-phosphate.
- 30 (24) 1-[4-(4-Chlorophenyl)butyl]-3-(2-phenylethyl)imidazolium chloride.

- (25) 1-[4-(4-Chlorophenyl)-1-methylbutyl]-3-heptylimidazolium
4-methylbenzenesulfonate.
- (26) 1-[4-(4-Chlorophenyl)-(E)-3-butenyl]-3-heptylimidazolium
chloride.
- 5 (27) 1-[4-(2,3-Dichlorophenyl)-2-butenyl]-3-heptylimidazolium
4-methylbenzenesulfonate.
- (28) 1-[4-(2-Chlorophenyl)-2-butenyl]-3-heptylimidazolium
dihydrogenphosphate.
- (29) 1-[4-(4-Chlorophenyl)-3-butynyl]-3-heptylimidazolium
4-methylbenzenesulfonate.
- 10 (30) 1,2-Dimethyl-3-[4-[4-[[methanesulfonyl]amino]phenyl]butyl]-
imidazolium methanesulfonate.
- (31) 1-[4-(2-Chlorophenyl)butyl]-3-heptylimidazolium dihydro-
genphosphate.
- 15 (32) 1-[4-(4-Chlorophenyl)butyl]-3-(2-methylpropyl)imidazolium
chloride.
- (33) 1-[4-(4-Chlorophenyl)butyl]-3-heptyl-2-methylimidazolium
dihydrogenphosphate.
- (34) 1-[4-(4-Chlorophenyl)butyl]-2,3-dimethylimidazolium chloride.
- 20 (35) 1-Heptyl-3-[4-(4-methoxyphenyl)butyl]-2,4,5-trimethyl-
imidazolium bromide.
- (36) 1-[4-(4-fluorophenyl)-2-oxobutyl]-3-heptylimidazolium
chloride.
- (37) 1-[5-(4-Chlorophenyl)pentyl]-3-heptylimidazolium dihydrogen-
phosphate.
- 25 (38) 1-[2-Hydroxy-2-[4-[(methanesulfonyl)amino]phenyl]ethyl]-3-
methylimidazolium iodide.
- (39) 1-Heptyl-3-[2-[4-[(methanesulfonyl)amino]phenyl]-2-oxoethyl]-
imidazolium bromide.
- 30 (40) 1-[2-((2,6-Dimethylphenyl)sulfamoyl)ethyl]-3-heptylimidazolium
bromide.


- (41) (E)-1-[4-Fluorophenyl]-3-butenyl]-3-heptylimidazolium dihydrogenphosphate.
- (42) (Z)-1-[4-(4-Chlorophenyl)-2-butenyl]-3-heptylimidazolium dihydrogenphosphate.
- 5 (43) 1-Methyl-3-[2-[4-((methylsulfonyl)-amino)phenyl]-ethyl]-1H-imidazolium iodide.
- (44) 1-Methyl-3-(2-[4-((methylsulfonyl)amino)phenyl]-2-oxoethyl)-1H-imidazolium bromide.
- 10 (45) 1-Methyl-3-[3-[4-((methylsulfonyl)-amino)phenyl]-3-oxopropyl]-1H-imidazolium chloride.
- (46) 1,2-Dimethyl-3-[2-[4-((methylsulfonyl)amino)phenyl]-2-oxoethyl]-1H-imidazolium chloride.
- (47) 1,2-Dimethyl-3-[2-hydroxy-2-[4-((methylsulfonyl)-amino)phenyl]ethyl]-1H-imidazolium chloride.
- 15 (48) 1-[2-[4-((Ethylsulfonyl)amino)-phenyl]-2-oxoethyl]-3-methyl-1H-imidazolium chloride.
- (49) 1-(1-Methylethyl)-3-[2-[4-((methylsulfonyl)amino)-phenyl]-2-oxoethyl]imidazolium chloride.
- 20 (50) 1-[2-Hydroxy-2-[4-((methylsulfonyl)amino)phenyl]ethyl]-3-(1-methylethyl)-1H-imidazolium chloride.
- (51) 1-[3-Hydroxy-3-[4-((methylsulfonyl)-amino)phenyl]-propyl]-3-methyl-1H-imidazolium chloride.
- (52) 1-Ethyl-3-[2-hydroxy-2-[4-((methylsulfonyl)amino)-phenyl]-1H-imidazolium chloride.
- 25 (53) 1-[2-[4-((Ethylsulfonyl)amino)phenyl]-2-hydroxyethyl]-3-methyl-1H-imidazolium chloride.

- (54) 1-Ethyl-3-[2-[4-((methylsulfonyl)-amino)phenyl]-2-oxoethyl]-1H-imidazolium chloride.
- (55) 1-Methyl-3-[3-[4-((methylsulfonyl)amino)phenyl]-propyl]-1H-imidazolium chloride.
- 5 (56) (-)-1-[2-Hydroxy-2-[4-((methylsulfonyl)amino)phenyl]-ethyl]-3-methyl-1H-imidazolium chloride.
- (57) (+)-1-[2-Hydroxy-2-[4-((methylsulfonyl)amino)phenyl]-ethyl]-3-methyl-1H-imidazolium chloride.

Among the compounds as defined by Formula I, a site for
 10 stereo or geometric isomerism may exist, *e.g.* asymmetric carbon atoms or olefinic linkages. Any of the optical and *cis/trans* isomers so possible are considered to be part of this invention.

Another aspect of this invention are the molecular compounds of the quaternary salts of Formula I with aromatic
 15 dicarboxylic acids. Representative of such acids are phthalic, acid, 1,8-naphthalenedicarboxylic acid, 1,2-naphthalenedicarboxylic acid and 2,3-naphthalenedicarboxylic acid.

A further aspect of this invention are certain intermediate substituted imidazoles which have been found to have
 20 antiarrhythmic activity. Such compounds are defined by the following Formula II:



II

wherein T is selected from the group consisting of nitro, amino, lower alkanesulfonamido, and Q is selected from the group consisting of a straight chain alkyl of 1-4 carbon atoms, $-\overset{\text{O}}{\underset{\text{O}}{\text{C}}}-\text{CH}_2-$, $-\text{CHOHCH}_2-$, and the pharmaceutically acceptable acid addition salts thereof. Representative of the compounds of Formula II are the following:

- (1) 1-[4-(4-Nitrophenyl)butyl]imidazole hydrochloride
- (2) N-[4-(1H-Imidaz-1-ylacetyl)phenyl]methanesulfonamide
- (3) 2-(1H-Imidaz-1-yl)-1-(4-nitrophenyl)ethanone hydrochloride
- 10 (4) N-[4-[1-Hydroxy-2-(1H-imidazol-1-yl)ethyl]phenyl]methanesulfonamide
- (5) 1-(4-Aminophenyl)-2-(1H-imidazol-1-yl)ethanone
- (6) α -(4-Nitrophenyl)-1H-imidazole-1-ethanol.
- (7) N-[4-[2--(1H-Imidazol-1-yl)ethyl]-phenyl]methanesulfonamide
- 15 (8) 1-[2-(4-Nitrophenyl)ethyl]-1H-imidazole hydrochloride
- (9) (+)-1-[2-Hydroxy-2-[4-((methylsulfonyl)amino)phenyl]ethyl]-1H-imidazole d-7,7-dimethyl-2-oxobicyclo[2.2.1]heptane-1-methane sulfonic acid salt
- 20 (10) (1)-1-[2-Hydroxy-2-[4-((methylsulfonyl)-amino)phenyl]ethyl]-1H-imidazole 1-7,7-dimethyl-2-oxo-bicyclo[2.2.1]-heptane-1-methanesulfonic acid salt
- (11) (-)-1-[2-Hydroxy-2-[4-((methyl-sulfonyl)amino)phenyl]ethyl]-1H-imidazole

In the foregoing Formula II lower alkyl defines a straight or branched chain of 1-4 carbon atoms. The pharmaceutically acceptable acid addition salts contemplated are prepared according to known procedures and are those derived from, for example, the following acids; hydrochloric, hydrobromic, sulfuric, phosphoric, acetic, propionic, benzoic, naphthoic, oxalic, succinic, maleic, malic, adipic, lactic, tartaric, citric, salicyclic, methanesulfonic and p-toluenesulfonic.

PROCESS ASPECT

The compounds of this invention may be prepared in general, by various processes and reactants known in the art. In order to produce the compounds according to Formula I and those intermediates of Formula II the following schemes were most usually employed.

In one schematic route, in order to produce a 1-substituted imidazole, combine one equivalent of an alkyl halide (or tosylate) or alternatively a substituted aralkyl halide (or tosylate) with 2-5 equivalents of imidazole. Heat the mixture neat or in a suitable solvent (e.g. acetonitrile, nitromethane, dimethylformamide) at from about 80 to about 150°C. Follow the progress of the reaction by thin-layer chromatography. When the reaction is completed, remove the solvent (if one used) *in vacuo* and dissolve the resultant reaction mixture in 5 to 10 volumes of water. If necessary, adjust the pH of the solution to pH 9 with aqueous base. Extract the aqueous mixture with methylene chloride, which is washed with water, dried over sodium sulfate and evaporated. In such a manner the 1-alkylimidazole or alternatively the 1-(substituted aralkyl)-imidazole may be prepared.

Still another method for preparing the foregoing compounds is as follows. To 1.1 equivalents of sodium hydride in anhydrous 1,2-dimethoxyethane with about 1.1 equivalents of dimethylformamide in an atmosphere of nitrogen add about one equivalent of imidazole. Heat the reaction mixture to reflux. When the

evolution of hydrogen has ceased, add the appropriate halide (or tosylate) slowly to the reaction mixture and when the addition is completed, continue heating and follow the progress of the reaction by thin-layer chromatography. When complete, quench the reaction mixture in ice water. Extract the aqueous mixture with methylene chloride, and as above, the appropriate 1-substituted imidazole is produced.

Generally, the 1,3-disubstituted imidazolium salts may be prepared as follows. Combine one equivalent of the 1-substituted imidazole with at least one or more equivalents of the alkyl halide (or tosylate) or substituted aralkyl halide (or tosylate). Heat the mixture neat or in a suitable solvent (e.g. acetonitrile, dimethylformamide or nitromethane) at about 80 to about 150°C. Follow the reaction by thin-layer chromatography and when complete, dissolve the reaction mixture (after removal of the solvent, if used, *in vacuo*) in water and adjust the pH of the solution with saturated sodium bicarbonate to about pH 8. Extract the aqueous solution several times with ether, then adjust the pH of the aqueous component with a suitable acid (same anion as that of the imidazolium salt) to about pH 3. Extract the aqueous layer with methylene chloride, after drying and evaporation of the solvent the appropriate imidazolium salt is produced.


It will be apparent to those skilled in the art that certain of the substituted aralkyl halides etc. are available commercially and still others may be prepared by variants on known processes.

One expeditious route to the substituted aralkyl halides especially in the case where the alkyl is an alkenyl moiety would be the diazotization of a suitably substituted aniline, and reacting

it to a Meerwein arylation with copper chloride, copper bromide or copper sulfate and a butadiene to produce the substituted arylbutenyl chloride, bromide or alcohol which latter compound can be converted to the tosylate ester.


The following schematics are illustrative of ways to produce the compounds of the invention:

A.




0131302





5

Meerwein
butadiene


10



15


 $H_2/Pd/C$ HBr or
PBr₃ $H_2/Pd/C$ 

20

CH₃NCO

25

RZ



30

If it is necessary or desirable to change the anion, this may be accomplished by commercially available ion exchange resins. As for instance, the chloride anion may be exchanged in an anion exchange resin (e.g. Biorad AG-1-X8, 20-50 mesh hydroxide form) and the resulting eluates titrated with 10% phosphoric acid to produce the $H_2PO_4^-$ anion.

USE AND PHARMACEUTICALCOMPOSITION ASPECT

5 The quaternized imidazoles of this invention, their
molecular compounds with aromatic dicarboxylic acids and certain
of the unquaternized imidazoles and their pharmaceutically
acceptable acid addition salts have been found to increase
refractoriness of cardiac tissue thereby providing their usefulness
as antiarrhythmic agents. Their activity has been analyzed
in several procedures such as, utilizing standard electro-
10 physiological techniques to measure resting potential, action
potential amplitude, duration, rate of rise and refractory
periods of normal canine Purkinje fibres; and also utilizing
the programmed electrical stimulation conscious dog model.

PHARMACOLOGICAL DATA

15 The primary method for evaluating these compounds
as antiarrhythmic agents was an in vitro electro-
physiological screen using canine Purkinje fibers.
Using conventional microelectrode techniques the effects
of the drugs on the action potential characteristics of
20 the Purkinje fiber were measured. Both Class I and
Class III antiarrhythmic agents can be distinguished
using this method. For a compound to be considered
active as a Class III agent it must have prolonged the
action potential duration at 95% repolarisation (APD_{95})
25 by 20% without affecting conduction properties. A
decrease in the rate of rise of phase 0 (V_{max}) by 20%
was considered a toxic effect on conduction for a
Class III agent as well as being the criterion for
activity for a Class I antiarrhythmic agent.

$C_{10}APD_{95}$ --the concentration of drug (in micromolar units) which causes a 10% increase or decrease (-) in APD_{95} from control value

5 $C_{20}APD_{95}$ --the concentration of drug (in micromolar units) which causes a 20% increase or decrease (-) in APD_{95} from control value

$MaxAPD_{95}$ (conc.)--the maximum observed effect on APD_{95} from control value and the concentration (in micromolar units) where this occurred

10 \dot{V}_{max} (conc.)--the change (+,-) in the rate of rise of phase 0 of the action potential from control value and the concentration of drug which caused the observed change

NR--not reached

15 NE--no effect

The above data were recorded for a normal stimulated beat at a basic cycle length of 1000 milliseconds.

20 The $C_{10}APD_{95}$ information is included here because some compounds which caused a 10% prolongation of APD_{95} but which did not cause a 20% prolongation in APD_{95} were shown to be effective in an in vivo dog model.

The following Table I illustrates the activity of some of the compounds.

TABLE I

In Vitro (Canine Purkinje Fiber Studies)

Compound No.	C ₁₀ APD ₉₅ (μM)	C ₂₀ APD ₉₅ (μM)	MaxAPD ₉₅ (conc)	V _{max} (conc)
I-1	(+) 12		27% (30)	-20% (15)
I-7	0.5	8	26% (10)	-12% (10)
I-12	20	NR		-20% (10-30)
I-14	13	30	21% (30)	increased
I-16	6.0	NR	11% (10)	-20% (>10)
I-19	1.0	10	20% (10)	-6% (10)
I-20	0.8	6.0	23% (10)	NE
I-21	2.0	6	25% (10)	NE
I-22	1.3	3.0	39% (10)	Minimal
I-25	10	NR	10% (10)	NE
I-29	20	NR	15% (30)	-20% (20)
I-30	1.0	7.0	23% (10)	NE
I-34	1.4	6.0	25% (10)	NE
I-38	0.3	0.8	63% (10)	-5% (10)
I-39	1.7	9.0	21% (10)	NE
I-42	38		12% (100)	-26% (100)
I-43	2.0	8.0	21% (10)	NE
I-44	1.0	2.3	38% (10)	NE
II-1	1.2	4.0	27% (10)	NE
II-2	3.0	14.0	67% (100)	-15% (100)
II-3	20.0	50.0	29% (100)	NE
II-4	2.0	7.0	23% (10)	-15% (10)
II-5	4.0	NR	13% (10)	Minimal
II-6	15.0	43.0	28% (100)	NE
II-7	2.0	10.0	47% (100)	NE
II-8	7.0	30.0	29% (100)	-15% (0.1)

Thus there is provided by this invention a method for treating arrhythmia which comprises administering to a subject suffering from an arrhythmia and in need of treatment or to a subject suspected of developing an arrhythmia an effective amount for treating such arrhythmia of a compound of this invention. The compounds are preferably utilized for the control of re-entrant arrhythmias in humans and for the prevention of sudden death resulting from ventricular fibrillation. Accordingly, it is contemplated that the compounds are best utilized in prophylactic treatment. Moreover, since the compounds enhance the electrical stability of the heart, they can be used in conjunction with electrical devices designed to terminate cardiac arrhythmias such as ventricular tachycardia and ventricular fibrillation.

In general, the compounds of this invention may be administered orally or parenterally. The dosage administered will be dependent on the subject being treated, the route of administration and the type and severity of the arrhythmia being prevented or reduced.

A typical dose for prophylactic treatment, however, will contain from about 0.5 mg/kg to about 5 mg/kg of the active compounds of this invention when administered orally. For I.V. administration, the dose will be from about 0.2 mg/kg to about 4 mg/kg, preferably about 0.2 to about 2 mg/kg.

The compound to be administered can be formulated by admixing with any of a number of suitable pharmaceutical diluents and carriers such as lactose, sucrose, starch powder, cellulose, calcium sulfate, sodium benzoate and the like. Such formulations can be compressed into tablets or can be encapsulated into gelatin capsules for convenient oral administration. Such a capsule may contain one of the compounds of this invention for example, 1-[4-(4-chlorophenyl)butyl]-3-heptylimidazolium dihydrogenphosphate or 2-(1H-imidaz-1-yl)-1-(4-nitrophenyl)-ethanone hydrochloride in the amount of about 10 to about 50 mg. Such formulation can be administered orally at the rate of about 1 or 2 capsules per day or more often as needed depending upon the particular condition and subject being treated.

For parenteral administration a compound of this invention can be formulated for intramuscular or intravenous administration. In the case of treatment of a patient suffering from a severe cardiac arrhythmia, it may be desirable to administer a compound of the invention by intravenous infusion in order to effect a rapid conversion to a normal sinus rhythm. The normalized condition can then be maintained by oral administration.

The compounds of this invention can be formulated for parenteral administration with any of a number of pharmaceutically acceptable carriers and diluents to constitute an injectable liquid solution. Commonly used diluents and carriers include water or saline solution, buffered aqueous solutions as well as dispersing and surface active agents if necessary. A typical formulation suited to intramuscular administration may contain one of the compounds of this invention such as 1-[4-(4-chlorophenyl)-2-

butenyl]-3-heptylimidazolium chloride (1:1) phthalic acid or
2-(1H-imidaz-1-yl)-1-(4-nitrophenyl)ethanone hydrochloride in the
amount of about 10 to 250 mg and a solubilizing agent and
sufficient sterile water to bring the volume to about 2 ml. Such
5 formulation can be injected at a rate of 1 to 4 times per day
or more often depending upon the particular condition of the
subject being treated.

The pharmaceutical preparations of the compounds of
this invention may optionally, additionally contain one or more
10 other pharmaceutically active substances. Some of the substances
envisioned are vasodilators such as glycerol trinitrate, pentaery-
thritol tetranitrate and carbochromene; diuretic agents, such
as chlorothiazide; heart tonics, such as digitalis preparations;
hypotensive agents, such as Rauwolfia alkaloids and guanethidine;
15 bronchodilators and sympathomimetic agents, such as isoprenaline,
orciprenaline, adrenalin and ephedrine; α -adrenergic blocking
agents, such as phentolamine; β -adrenergic blocking agents, such
as propranolol and other antiarrhythmic agents such as quinidine.

Other utilities may be ascribed to certain of the
20 quaternized imidazoles of this invention, e.g. cardiotonic agents,
bradycardic agents, antifungal agents and stabilizers for
physiological formulations such as ophthalmic drops.

This invention described hereinabove is illustrated
below in the Examples, which, however, are not to be construed
25 as limiting the scope of this invention.

PREPARATION 1Acetic Acid, 4-[4-(Acetylamino)phenyl]butyl ester

To a solution of 30.0 g (0.15 mole) of 4-(p-nitrophenyl)-
butanol in 375 ml glacial acetic acid add 30.9 g (0.55 mole) of
5 iron filings. Stir the suspension at reflux for 5 1/2 hrs,
following the progress of the reaction by thin-layer chromatography
on silica gel (acetonitrile: ammonium hydroxide, 90:10). At the
completion of the reaction, pour the reaction mixture onto 3 liters
of ice water and extract with 3 x 300 ml of chloroform. Combine
10 the organic layers and wash with 100 ml water. Dry the chloroform
extracts over potassium carbonate, filter and concentrate *in vacuo*
to give an oily solid. Recrystallize from 400 ml ether to obtain
the title compound.

NMR (CDCl₃): δ =1.50-1.85(m,4), 2.04(s,3), 2.14(s,3),
15 2.60(t,2), 4.06(t,2), 6.92-7.55(m,4)
and 7.90(br s,1)ppm.

PREPARATION 24-(Acetylamino)benzenebutanol

To a solution of 19.3 g (0.077 mole) of acetic acid,
20 4-[4-(acetylamino)phenyl]butyl ester in 300 ml of ethanol, add a
solution of 21 g potassium carbonate in 300 ml of water. Heat
the resulting solution at 55°C for 4 hrs, following the progress
of the reaction by thin-layer chromatography on silica gel
(acetonitrile: ammonium hydroxide, 90:10). At the completion
25 of the reaction, filter off any insolubles and remove the ethanol
in vacuo. Extract the aqueous residue with 3 x 300 ml methylene
chloride. Wash the combined organic layers with 100 ml water,
dry over potassium carbonate, filter and remove the solvent
in vacuo. Recrystallize the resulting solid from 140 ml acetonitrile
30 to obtain the title compound.

NMR (CDCl₃): δ =1.50-1.90(m,4), 1.75(s,2, exchange),
2.15(s,3), 2.45-2.80(m,2), 3.65(t,2)
and 7.00-7.60(m,4)ppm.

PREPARATION 34-Methylbenzenesulfonic acid, 4-[4-(acetylamino)phenyl]butyl ester

To a 0°C solution of 4.5 g (0.022 mole) 4-(acetylamino)-benzenebutanol in 25 ml pyridine, add 8.3 g (0.044 mole) tosyl chloride and chill in the refrigerator overnight. Follow the progress of the reaction by thin-layer chromatography on silica gel (acetonitrile: ammonium hydroxide, 90:10). At the completion of the reaction, pour the reaction mixture onto 500 ml ice water and stir for 15 minutes. Extract the aqueous mixture with 3 x 300ml ether. Wash the combined organic layers with 1N hydrochloric acid, then with water, dry over potassium carbonate, filter and remove the solvent *in vacuo* to obtain the title compound.

NMR (CDCl₃): δ=1.40-2.00(m,4), 2.10(s,3), 2.40(s,3), 2.30-2.80(m,2), 4.00(t,2) and 6.90-7.90(m,8)ppm.

PREPARATION 41-Heptyl-2,4,5-trimethyl-1H-imidazole

Dissolve 16.0 g (0.019 mole) of 2,4,5-trimethyl-1H-imidazole hydrochloride in 125 ml dry N,N-dimethylformamide and cool to 0°C. Add 8.75 g (0.218 mole) of sodium hydride (60%), stir at 0°C for 30 minutes and then heat reaction mixture to 100°C for one hour. Cool to 0°C and then add 18.0 ml (0.114 mole) of n-heptyl bromide to reaction mixture. Upon completion of addition, heat to 100°C for four hours, following the progress of the reaction by thin-layer chromatography on silica gel (methanol: 1M sodium chloride, 95:5). At the completion of the reaction, cool to room temperature and add 2 ml water. Filter precipitated salts and evaporate solvent.

Dissolve this residue in 200 ml 10% NaOH and then extract into methylene chloride (200 ml). Evaporate solvent and then distill *in vacuo* to obtain the title compound.

NMR (CDCl₃): δ=0.90(t,3), 1.1-1.7(m,10), 2.1(s,6), 2.4(s,3) and 3.76(t,2)ppm.

PREPARATION 51-Chloro-4-ethynylbenzene

Dissolve 276.5 g (0.634 mole) of bromomethyltriphenylphosphonium bromide in 1800 ml THF and cool to -78°C . Add 67.6 g (0.062 mole) potassium tert.-butoxide and stir at -78°C for 2 hours. Add 89.12 g p-chlorobenzaldehyde in 200 ml THF to cold reaction mixture and stir for 30 minutes. Add 142.3 g potassium tert.-butoxide and then allow to warm to room temperature. After 18 hours, add 500 ml saturated aqueous ammonium chloride solution and 500 ml H_2O and extract with 500 ml diethyl ether. Wash organic layer with 2 x 400 ml saturated aqueous sodium chloride solution, dry over magnesium sulfate, filter and evaporate solvent to dryness. Wash residue with 200 ml hexane and filter off precipitate. Evaporate hexane and dissolve material in 150 ml hexane:methylene chloride (66:33) solution and elute through column containing 300 g silica gel (Merck 60-230 mesh) with hexane:methylene chloride (66:33) solution. Monitor column chromatography using thin-layer chromatography on silica gel (hexane:methylene chloride, 66:33). Collect desired fraction and evaporate solvent. Distill residue *in vacuo* and collect fraction boiling at 65°C (8.5mmHg) to obtain the title compound.

PREPARATION 64-(4-Chlorophenyl)-3-butyn-1-ol

Dissolve 14.2 g (0.104 mole) of 1-chloro-4-ethynylbenzene in 35 ml dry THF and cool to -78°C . Add 49 ml butyllithium to cold reaction mixture and allow to warm to room temperature for 2 hours. After 2 hours, cool to -78°C and then add 5.8 ml (0.114 mole) ethylene oxide in 20 ml THF. Allow to warm to room temperature. Monitor progress of reaction with thin-layer chromatography on silica gel (methylene chloride).

After 18 hours, add 150 ml saturated aqueous ammonium chloride solution and extract with 150 ml diethyl ether. Wash organic layer with 2 x 100 ml saturated aqueous sodium chloride solution. Dry organic layer with magnesium sulfate and then filter, and evaporate solvent to obtain the title compound.

PREPARATION 7

4-(4-Chlorophenyl)-3-butyne-1-ol 4-methylbenzenesulfonate (Ester)

Dissolve 13.3 g (0.0736 mole) of 4-(4-chlorophenyl)-3-butyne-1-ol in 70 ml pyridine and cool to 0°C. Add 16 g (0.0837 mole) of p-toluenesulfonyl chloride and stir in ice bath for one hour. Store in freezer over night. Add 200 ml diethyl ether and then wash with 3 x 150 ml 1N hydrochloric acid. The organic layer is then washed with 50 ml saturated aqueous sodium bicarbonate solution and then dried over magnesium sulfate. Evaporate solvent to obtain the title compound.

NMR (CDCl₃): δ =2.42(s,3), 2.77(t,2), 4.17(t,2), 7.21(s,4), 7.31(d,2) and 7.82(d,2)ppm.

PREPARATION 8

4-(Methanesulfonylamino)benzenebutanol methanesulfonate (Ester)

9.25 g (0.056 mole) 4-aminobenzenebutanol is dissolved in 100 ml pyridine and the solution is cooled to ca. -10°C. 17.33 ml (25.65 g) (0.244 mole) methanesulfonyl chloride is added dropwise with vigorous stirring at a rate such that the reaction temperature does not rise beyond -5°C. After all methanesulfonyl chloride is added, the reaction mixture is left in the freezer for two days. The reaction mixture is poured onto 300 ml ice water and extracted with 3 x 200 ml methylene chloride. The combined organic layers are washed with 1 liter cold 5% HCl in 3 portions. The organic layer is dried over sodium sulfate, filtered, and stripped. The residue is crystallized from ethyl acetate/ether to obtain the title compound.

NMR (CDCl_3): δ =1.56-1.93 (m, 4), 2.47-2.82 (m, 2), 2.98 (s, 3), 3.02 (s, 3), 4.07-4.42 (m, 2), 7.20 (s, 4) and 7.25 (br s, 1) ppm.

PREPARATION 9

5 2,3-Dichlorophenyldiazonium chloride

To a suspension of 25 g (0.15 mole) 2,3-dichloroaniline in 16 ml H_2O , add 39 ml (0.46 mole) concentrated HCl. Stir the mixture at room temperature for about five minutes. Cool the mixture to -5 to 0°C and add dropwise a solution of 11 g (0.16 mole) NaNO_2 in 30 ml H_2O . Stir the resulting solution at -5 to 0°C for about 30 minutes, then filter the solution into an ice-cooled filter flask. Keep the filtrate containing the diazonium salt at about 0°C.

PREPARATION 10

15 1-Chloro-4-(2,3-dichlorophenyl)-2-butene

Suspend 12.3 g (0.09 mole) $\text{NaOAc} \cdot 3\text{H}_2\text{O}$ and 4.6 g (0.03 mole) CuCl_2 in 160 ml acetone and 25 ml H_2O . Cool the mixture to -5 to 0°C. Add 12.4 g (20 ml, 0.23 mole) of condensed butadiene. Add the solution of 2,3-dichlorophenyldiazonium chloride dropwise, keeping the diazonium salt solution and the reaction mixture between -5 and 5°C at all times. When the addition is complete, discontinue cooling, and stir the reaction mixture at room temperature for about 16 hours.

20 Add 160 ml diethyl ether to the reaction mixture and stir at room temperature for about five minutes. Separate the layers, and wash the organic layer with 2 x 150 ml H_2O and 150 ml saturated NaCl solution. Dry the organic layer over MgSO_4 . Evaporate the solvent, and distill the resulting oil to obtain the title compound.

30 NMR (CDCl_3): δ =3.52 (d, 2), 4.05 (d, 2), 5.01-6.27 (m, 2) and 7.04-7.52 (m, 3) ppm.

PREPARATION 11Cyclopropyl-4-fluorophenylmethanone

Add 20.0 g (0.10 mole) γ -chloro-p-fluorobutyrophenone dropwise to a methanolic KOH solution (prepared from 9.8 g of 86% KOH pellets and 60 ml of methanol). Stir the mixture at room temperature for 40 minutes and pour into 100 ml of water and then extract with 3 x 30 ml of methylene chloride. Wash the methylene chloride extracts with 3 x 40 ml of water, dry over magnesium sulfate and evaporate off the solvent to provide the title compound.

PREPARATION 12 α -Cyclopropyl-4-fluorobenzenemethanol

Add 3.2 g (0.085 mole) cyclopropyl-4-fluorobenzylmethanone to 80 ml of 3.2 g (0.085 mole) sodium borohydride partially dissolved in absolute ethanol, following the reaction by thin-layer chromatography. After about 6 hours at room temperature, cool the reaction in an ice bath and add acetic acid. Extract the mixture with methylene (3 x 60 ml) wash the methylene chloride extracts with saturated aqueous sodium bicarbonate solution, and dry over magnesium sulfate. Evaporate the methylene chloride extracts and dry *in vacuo* to yield the title compound.

PREPARATION 13(E)-1-(4-Chloro-1-butenyl)-4-fluorobenzene

Dissolve 12.8 g (0.077 mole) of α -cyclopropyl-4-fluorobenzenemethanol in 60 ml of CHCl_3 , add 20 ml of concentrated HCl and reflux overnight. Separate the two layers, and extract the aqueous layer with methylene chloride. Combine the methylene chloride extracts with the chloroform layer and wash with saturated aqueous sodium bicarbonate, and dry over magnesium sulfate. Evaporate the organic solvents to yield the title compound.

PREPARATION 141-Chloro-4-(2-propynyl)benzene

In a flask containing 59 g (2.43 mole) magnesium turnings in 500 ml anhydrous ether under an atmosphere of nitrogen add slowly a solution of 465 g (2.43 mole) of 4-bromochlorobenzene in 1 liter 4 anhydrous ether. With caution, once the Grignard has started maintain slow reflux until all 4-bromochlorobenzene has been added. Stir an additional hour and then add this Grignard (maintain slow reflux) to a suspension of 170 g (2.43 mole) methoxyallene and 69 g of cuprous bromide in 300 ml anhydrous ether under N₂. When addition is complete, stir another 10 minutes and then add saturated ammonium chloride. Decant off the ether solution wash the solid residue with ether (3 x 50 ml). Combine the ether extracts and wash with 1N HCl, saturated aqueous sodium bicarbonate, water and dry over magnesium sulfate. Evaporate off the ether and distill the residue *in vacuo* to obtain the title compound (99-105mmHg).

PREPARATION 151-(4-Chlorophenyl)-2-butyne-1-ol

Under nitrogen and with cooling, to a solution of 150.6 g (1.0 mole) 1-chloro-4-(2-propynyl)benzene in 600 ml of anhydrous ether, add dropwise 400 ml of 2.5M *n*-butyllithium (1.0 mole) in hexane. When addition is complete, stir a further 30 minutes and under N₂ add 31 g of para-formaldehyde. Stir the mixture in a dry ice/acetone bath for 1 hour. Allow to warm to room temperature and then allow to reflux. When the reaction is complete, add 400 ml of saturated aqueous sodium chloride. Separate the layers and extract the aqueous layer with ether. Wash the combined ether solution, wash with water and dry over magnesium sulfate. Evaporate off the ether, dissolve the

residue in hot cyclohexane which yielded crystalline material.
the title compound.

PREPARATION 16

(Z)-1-(4-Chlorophenyl)-2-buten-1-ol

5 Under nitrogen, hydrogenate a mixture of 10 g (0.055 mole) of 1-(4-chlorophenyl)-2-butyne-1-ol and 0.6 g Lindlar catalyst. Following the reaction *via* NMR the reaction is complete in about two days. The hydrogenation is stopped and the reaction stirred under N₂ at room temperature overnight. Filter off the
10 catalysts, wash with methylene chloride combine the filtrate and evaporate to yield the title compound.

PREPARATION 17

(Z)-4-(4-Chlorophenyl)-2-buten-1-ol 4-methylbenzenesulfonate

Dissolve 6.7 g (0.036 mole) of (Z)-1-(4-chlorophenyl)-
15 2-buten-1-ol in 100 ml of anhydrous ether, add 3.2 g of powdered KOH. Stir the mixture and with cooling add a solution of 7.0 g p-toluenesulfonylchloride in 30 ml of anhydrous ether. Continue to stir with cooling for 30 minutes, then allow to warm to room temperature overnight. Filter, wash the filtrate three
20 times with saturated aqueous sodium bicarbonate, then with water and dry over sodium sulfate. Evaporate off solvents and triturate residue with petroleum ether, cyclohexane. Recrystallize residue from cyclohexane to yield the title compound.

PREPARATION 18N-Phenylmethanesulfonamide

To a 0°C solution of 250 g (2.68 mole) of aniline and 233.2 g (2.95 mole) of pyridine in 1.25 L of methylene chloride and 319.6 g (2.79 mole) of methanesulfonyl chloride dropwise. Stir the resulting solution at 0°C for about 30 minutes then warm to ambient temperature for about 16 hours. Extract the mixture with 2N aqueous sodium hydroxide (4x1 L). Combine the aqueous extracts and wash with 1 L of methylene chloride. Cool the aqueous layer to about 0°C and acidify with concentrated hydrochloric acid until about pH 1. Collect the precipitate and dry to give the title compound.

NMR (DMSO-d₆): δ = 2.97(s,3), 7.10(t,1),
7.20(d,2), 7.33(t,2), and
9.72(br s, 1)ppm.

PREPARATION 19N-Phenylethanesulfonamide

In a manner similar to PREPARATION 18 utilize aniline + ethanesulfonyl chloride to prepare the title compound.

PREPARATION 20N-[4-(2-Chloro-1-oxoethyl)phenyl]methanesulfonamide

5 In a flask containing 128.4 g (0.75 mole) of N-phenylmethanesulfonamide, 169.5 g (1.50 mole) of chloro-
acetyl chloride in 1 L of methylene chloride under an
atmosphere of nitrogen and cooled to about -10°C add
300 g (2.25 mole) of aluminum chloride. Stir the
resulting mixture for about 2 hours at -10°C then warm to
room temperature for about 16 hours. Pour the reaction
10 mixture onto 2 Kg of ice containing 600 mL of
concentrated hydrochloric acid. Collect the precipitate
and wash with methanol (3x300 mL) and dry in vacuo
to give the title compound.

15 NMR (DMSO-d₆): δ = 3.16(s,3), 5.15(s,2),
7.40(d,2), 8.06(d,2),
and 10.48 (br s, 1)ppm.

PREPARATION 21N-[4-(3-Chloro-1-oxopropyl)phenyl]methanesulfonamide

20 In a manner similar to PREPARATION 20 utilizing
N-phenylmethanesulfonamide + chloropropionyl chloride
prepare the title compound.

EXAMPLE 1

1-[4-(4-Chlorophenyl)butyl]-3-heptylimidazolium chloride (1:1)
compound with phthalic acid

5 1-Heptyl-1H-imidazole (6.67 g, 0.04 mole) and 4-(4-chlorophenyl)butyl chloride are heated and stirred at ca. 130°C. Follow the progress of the reaction by thin-layer chromatography on silica gel (acetonitrile: ammonium hydroxide, 9:1). At the completion of the reaction, the mixture is cooled to 20-25°C and triturated with diethyl ether to remove any untreated starting
10 material. The residue is dissolved in 100 ml of acetone. To the acetone solution of imidazolium salt is added a solution of phthalic acid (7.52 g, 0.045 mole) in 50 ml of acetone. The mixture is warmed to effect complete solution, then the solvent is removed *in vacuo*. The residue is recrystallized from acetone
15 to provide the title compound.

NMR (DMSO-d₆): δ=0.56-2.18(m,17), 2.64(t,2), 4.00-4.53
(m,4), ca. 5.0-ca. 9.0(broad,2), 7.09-
7.50(m,4), 7.50-7.90(m,4), 7.99(m,2)
and 9.70(m,1)ppm.

EXAMPLE 2

1-[3-(4-Methoxyphenyl)propyl]-3-methylimidazolium bromide

A mixture of 2.50 g (0.03 mole) of 1-methyl-1H-imidazole and 6.9 g (0.03 mole) of 1-(3-bromopropyl)-4-methoxy-
25 benzene is heated at 140°C for 1.5 hours. Follow the progress of the reaction by thin-layer chromatography on silica gel (methanol: 1M sodium chloride, 95:5). At the completion, the cooled reaction product is triturated thoroughly with ether and on drying, provides
30 the title compound.

NMR (CDCl₃): δ=2-2.09(m,4.5), 3.7(s,3), 4.0(s,3), 4.07-
4.08(t,2), 6.6-7.2(quar.,4), 7.4-7.7
(m,2) and 10.15(s,1)ppm.

EXAMPLE 31-Butyl-3-[3-(4-methoxyphenyl)propyl]imidazolium bromide

A mixture of 2.5 g (0.02 mole) of 1-butyl-1*H*-imidazole and 4.61 g (0.02 mole) of 1-(3-bromopropyl)-4-methoxybenzene is heated at 140°C for about 1.5 hours. Follow the progress of the reaction by thin-layer chromatography on silica gel (methanol: 1*M* sodium chloride, 95:5). At the completion of the reaction, the cooled reaction product is triturated thoroughly with ether and on drying, provides the title compound.

NMR(CDCl₃): δ=1.0-3.0(m,11), 3.3-4.0(m,5), 4.2-4.8(m,4), 6.75-7.45(m,4), 7.7-7.9(bs,2) and 10.4 (s,1)ppm.

EXAMPLE 41-Heptyl-3-[3-(4-methoxyphenyl)propyl]imidazolium bromide

A mixture of 2.5 g (0.015 mole) of 1-heptyl-1*H*-imidazole and 3.45 g (0.015 mole) of 1-(3-bromopropyl)-4-methoxybenzene is heated at 140°C for about 1.5 hours. Follow the progress of the reaction by thin-layer chromatography on silica gel (methanol: 1*M* sodium chloride, 95:5). At the completion of the reaction, the cooled reaction product is triturated thoroughly with ether and on drying, provides the title compound.

NMR(CDCl₃): δ=0.8-2.8(m,19), 3.72(s,31), 4.15-4.6(m,4), 6.7-7.2(m,4), 7.5-7.75(m,2) and 10.4 (s,1)ppm.

EXAMPLE 51-[4-(4-Fluorophenyl)-4-oxobutyl]-3-heptylimidazolium bromide

To 3.83 g (0.019 mole) γ-chloro-p-fluorobutyrophenone add 3.19 g (0.019 mole) 1-heptyl-1*H*-imidazole and lower the mixture into a constant temperature bath set for 140°C. Stir the mixture for about 26 hours at this temperature. Follow the reaction progress by thin-layer chromatography on silica gel

(methanol: 1M sodium chloride 95:5). At the completion of the reaction, triturate the cooled oil with ethyl acetate to obtain a white crystalline solid. Dry *in vacuo* to obtain the title compound.

NMR(CDCl₃): δ =0.65-1.5(br m) + 1.5-3.2(br m, 21), 3.8-4.1(t, 2), 4.2-4.7(m, 2), 6.9-7.4(br m, 5) and 7.4-7.8(br m, 2)ppm.

EXAMPLE 6

1-Heptyl-3-[4-(4-methoxyphenyl)butyl]imidazolium bromide

A mixture of 2.5 g (0.015 mole) of 1-heptyl-1H-imidazole and 3.66 g (0.015 mole) of 1-(4-bromobutyl)-4-methoxybenzene is heated at 140°C for about 2 hours. Follow the progress of the reaction by thin-layer chromatography on silica gel (methanol: 1M sodium chloride, 95:5). At the completion of the reaction, the cooled reaction product is triturated thoroughly with ether and on drying, provides the title compound.

NMR(CDCl₃): δ =0.6-2.2(m, 18), 2.3-2.75(br t, 2), 2.85-3.2(br s, 1), 3.7(s, 3), 4.05-4.48(m, 4), 6.6-7.2(m, 4), 7.52(br s, 2) and 10.3(s, 1)ppm

EXAMPLE 7

1-Heptyl-3-[4-(4-methoxyphenyl)butyl]-2-methylimidazolium bromide

A mixture of 2.50 g (0.0139 mole) of 1-heptyl-2-methyl-1H-imidazole and 3.37 g (0.0139 mole) of 1-(4-bromobutyl)-4-methoxybenzene is heated at 140°C for about 2 hours. Follow the progress of the reaction by thin-layer chromatography on silica gel (methanol: 1M sodium chloride, 95:5). At the completion of the reaction, the cooled reaction product is triturated thoroughly with ether and on drying, provides the title compound.

NMR(CDCl₃): δ =0.80-2.00(m, 17), 2.50-2.70(t, 2), 2.80(s, 3), 3.80(s, 3), 4.10-4.45(m, 4), 6.70-7.25(m, 4) and 7.70(s, 2)ppm.

EXAMPLE 81-[3-((2,6-Dimethylphenyl)sulfamoyl)propyl]-3-heptylimidazolium chloride

Combine 15.0 g (0.0919 mole) 1-heptyl-1*H*-imidazole
5 and 25.5 g (0.0974 mole) 3-chloro-*N*-(2,6-dimethylphenyl)propane-
sulfonamide and heat for about 24 hours at 130°C. Follow the
progress of the reaction by thin-layer chromatography on silica
gel (acetonitrile: ammonium hydroxide, 9:1). At the completion
of the reaction, dissolve the resulting cooled oil in 500 ml H₂O.
10 Extract the resultant solution with 6 x 500 ml hexane. Extract
aqueous layer with 3 x 500 ml CH₂Cl₂. Wash combined CH₂Cl₂ layers
1 x 250 ml saturated NaCl solution. Dry the methylene chloride
solution over Na₂SO₄. Evaporate the CH₂Cl₂ to provide the title
compound as an oil. Triturate the oil with Et₂O to provide solid.
15 Recrystallize by dissolving the solid in a minimum of hot acetone.
Seed and chill the solution to provide white crystals.

NMR(CDCl₃): δ=0.90-1.50(m,11), 1.5-2.1(m,2), 2.4(s,6),
2.4-3.0(m,2), 3.0-3.7(m,2), 3.9-4.3(t,2),
4.3-4.9(m,2), 7.0(s,3), 7.2(s,1), 7.89
20 (s,1), 9.15(s,1) and 10.0(s,1)ppm.

EXAMPLE 91-[4-(2,3-Dichlorophenyl)butyl]-3-heptylimidazolium chloride

A solution of 17 g (0.042 mole) 1-[4-(2,3-dichloro-
phenyl)-2-butenyl]-3-heptylimidazolium chloride in 250 ml
25 absolute ethanol and 1.5 g 10% Pd/C catalyst are placed in a small
Parr bottle, and hydrogenated at 25 psi for ca. 2 hours.
Additional H₂ is added as needed to maintain 25 psi pressure.
The reaction mixture is filtered and the solvent is evaporated
to provide the title compound.

30 NMR(CDCl₃): δ=0.45-1.56(m,1), 1.56-2.34(m,6), 2.86(t,2),
4.09-4.67(m,4), 6.90-7.48(m,3), 7.58(s,1),
7.74(s,1) and 10.75(s,1)ppm.

EXAMPLE 10

1-Heptyl-3-(4-phenylbutyl)imidazolium chloride (1:1 compound with phthalic acid)

Combine 5.35 g (0.32 mole) of (4-chlorobutyl)benzene and 5.27 g (0.32 mole) 1-heptyl-1H-imidazole, stir the mixture at approximately 145°C for about 30 hours. Follow the progress of the reaction by thin-layer chromatography on silica gel (acetonitrile: ammonium hydroxide 9:1). At the completion of the reaction, dissolve the resulting cooled oil in methanol and evaporate the solvent. Dissolve this product in 125 ml H₂O and extract the solution with 5 x 100 ml hexane. Then extract the aqueous with 2 x 50 ml methylene chloride. Combine the methylene chloride layers and wash with 1 x 100 ml saturated aqueous NaCl solution. Dry the methylene chloride extracts over sodium sulfate. Evaporate the solvents to obtain 1-heptyl-3-(4-phenylbutyl)imidazolium chloride as an oil. Dissolve the oil in approximately 35 ml acetone, and to it add an equimolar amount of phthalic acid. Remove the solvents *in vacuo* to obtain the title compound, which may be recrystallized from acetone.

NMR(DMSO-d₆): δ=0.55-1.50(m,13), 1.50-2.20(m,4), 2.42-2.90(t,2), 4.00-4.55(m,4), 7.25(s,5), 7.50-7.80(m,4), 7.91(m,2), 9.65(s,1) and 10.0-13.0(bs,2)ppm.

EXAMPLE 11

1-[4-(4-Chlorophenyl)butyl]-3-heptylimidazolium dihydrogenphosphate

Dissolve 31 g (0.084 mole) of 1-[4-(4-chlorophenyl)-butyl]-3-heptylimidazolium chloride in 70 ml of water and add to a 450 g of anion exchange resin column (Biorad AG 1-X8, hydroxide form). Elute with deionized water until the pH of eluent is about 8. Extract the combined eluent with 2 x 250 ml of diethyl ether. Adjust the pH of the aqueous solution to about 4 with phosphoric acid.

Remove the water *in vacuo*, recrystallize the solid residue from acetonitrile, then from 2-propanol to provide the title compound.

5 NMR (CD_3OD): δ =0.80-2.22(m,17), 2.70(t,2), 4.11-4.50 (m,4), 7.24(s,4), 7.62-7.75(m,2) and 9.17(bs,1)ppm.

EXAMPLE 12

1-[4-[4-(Acetylamino)phenyl]butyl]-3-heptylimidazolium 4-methylbenzenesulfonate

10 Heat a mixture of 6.23 g (0.017 mole) 4-methylbenzenesulfonic acid 4-(4-acetylamino)butyl ester and 2.9 g (0.017 mole) 1-heptyl-1*H*-imidazole to 80°C, for 5 hours, following the reaction by thin-layer chromatography on silica gel (acetonitrile: ammonium hydroxide, 90:10). At the completion of the reaction, 15 triturate the cooled resultant material with petroleum ether. Recrystallize the material from 25 ml hot methyl ethyl ketone with hot filtration, chill and seed to obtain the title compound.

20 NMR ($\text{DMSO}-d_6$): δ =0.60-1.00(t,3), 1.00-2.00(br m,14), 2.25(s,3), 2.30(s,3), 2.35-2.70(t,2), 3.94-4.35(m,4), 6.90-7.90(m,10), 9.25 (s,1) and 9.85(s,1)ppm.

EXAMPLE 13

1-[3-(4-Chlorophenyl)propyl]-3-heptylimidazolium bromide

25 A mixture of 2.0 g (0.012 mole) of 1-heptyl-1*H*-imidazole and 3.0 g (0.012 mole) of 1-(3-bromopropyl)-3-chlorobenzene is heated at 115°C for 4 hours, following the progress of the reaction by thin-layer chromatography on silica gel (methanol: 1*M* sodium chloride 95:5). At the completion of the reaction, the reaction mixture is dissolved in water and methanol mixture and it is then 30 washed with cyclohexane. The title compound is then extracted out of the aqueous solution with methylene chloride and obtained by removal of the methylene chloride as the hemihydrate.

NMR (CDCl₃): δ =0.60-1.55(m,11), 1.62-2.10(m,2), 2.60(m,2), 2.60-3.10(m,2), 4.20-4.55(m,4), 7.20(s,4), 7.42(bs,1), 7.70(bs,1) and 10.30(bs,1)ppm.

5

EXAMPLE 14

1-[4-(4-Chlorophenyl)butyl]-3-heptyl-2-methylimidazolium chloride (1:1) compound with phthalic acid

To 1.60 g (0.0078 mole) 1-chloro-4-(4-chlorobutyl)-benzene add 1.40 g (0.0078 mole) 1-heptyl-2-methyl-1H-imidazole. Stir the solution for one hour at about 120°C. Cool and triturate the oil with 10 ml EtOAc, followed by 2 x 10 ml petroleum ether, followed by 10 ml diethyl ether. Dissolve the oil in MeOH, and elute through a 1" bed of flash chromatography silica gel (Baker) with MeOH. The solvent is removed to yield an oil. Add to 0.64 g of the oil, 10 ml acetone and 0.51 g (0.0031 mole) phthalic acid. Remove the acetone *in vacuo*. Recrystallize the residue from acetone, filter and dry *in vacuo* to yield the title compound as a 1.25 hydrate.

NMR (CDCl₃+DMSO-d₆): δ =0.73-2.17(br m,17), 2.73(s,3), 4.00-4.47(br m,4), 7.25(s,4) and 7.40-7.87(br m,6)ppm.

20

EXAMPLE 15

1-[4-(4-Chlorophenyl)butyl]-2,3-dimethylimidazolium chloride

To 4.77 g (0.023 mole) 1-chloro-4-(4-chlorobutyl)-benzene add 2.23 g (0.023 mole) 1,2-dimethyl-1H-imidazole. Stir the solution for three hours at about 115°C. Follow the progress of the reaction by thin-layer chromatography on silica gel (MeOH: NaCl(1M), 92:8). At the completion of the reaction, cool to obtain an oil. Triturate the oil with petroleum ether until a solid precipitates. Recrystallize the solid from MeOH/acetone; filter and dry *in vacuo* to yield the title compound as a monohydrate.

30

NMR (CDCl₃+DMSO-d₆): δ =1.43-1.93(br m,4), 2.43-2.83 (br m,2), 2.67(s,3), 3.83(s,3), 4.22(t,2), 7.23(s,4) and 7.72 (s,2)ppm; H₂O at 3.3ppm.

35

EXAMPLE 161-[2-[(2,6-Dimethylphenyl)sulfamoyl]ethyl]-3-heptylimidazolium
bromide

Combine 3.46 g (0.019 mole) 1-bromoheptane and 5.16 g
5 (0.018 mole) N-(2,6-dimethylphenyl)-2-[1H-imidazol-1-yl]ethane
sulfonamide in 10 ml DMF and heat to 100°C. Follow progress of
reaction by TLC on silica gel (acetonitrile: ammonium hydroxide,
9:1). After about 10 hours, allow reaction mixture to cool to
room temperature. Pour the reaction mixture into 50 ml H₂O and
10 extract with 1 x 25 ml ether to give a white precipitate in the
aqueous layer. Filter and recrystallize the precipitate from
acetone to provide the title compound.

NMR: (DMSO-d₆): δ=0.60-1.55(m,11), 1.55-2.05(m,2),
2.40(s,6), 3.70-4.05(t,2), 7.18(s,3),
15 7.85-8.10(m,2), 9.20-9.40(bs,1) and
9.50(s,1)ppm.

EXAMPLE 17(E)-1-[4-(4-Chlorophenyl)-2-butenyl]-3-heptylimidazolium chloride
(1:1) compound with phthalic acid

20 A mixture of 1.65 g (0.01 mole) of 1-heptyl-1H-imidazole
and 2.00 g (0.01 mole) of (E)-1-chloro-4-(4-chlorophenyl)-2-butene
is heated at 105°C for about two hours, following the progress of
the reaction by thin-layer chromatography on silica gel (methanol:
1M sodium chloride, 95:5). At the completion of the reaction, the
25 reaction mixture is dissolved in water and it is then washed with
cyclohexane. The imidazolium salt is then extracted out of the
aqueous solution with methylene chloride and obtained by removing
the organic solvent.

A solution of 0.39 g (2.34 mmole) of phthalic acid and
30 0.86 g (2.34 mmole) of (E)-1-[4-(4-chlorophenyl)-2-butenyl]-3-heptyl-
imidazolium chloride in acetone is evaporated and then triturated
with ether, filtered to obtain the title compound.

NMR (CDCl_3): δ =0.55-2.15(m,13), 3.20-3.45(d,2), 4.15-4.55(t,2), 4.90-5.10(d,2), 5.70-6.25(m,2), 6.90-7.40(m,10) and 10.30(s,2)ppm.

EXAMPLE 18

5 A. 1-[4-(4-Chlorophenyl)butyl]-3-propylimidazolium chloride

Add 7.13 g (0.035 mole) of 1-chloro-4-(4-chlorobutyl)-benzene to 3.52 g (0.032 mole) of 1-propyl-1H-imidazole and heat to 130°C for 18 hours. Follow the progress of reaction by NMR (CDCl_3). At the completion of the reaction, cool to room temperature to obtain the title compound.

10 B. 1-[4-(4-Chlorophenyl)butyl]-3-propylimidazolium dihydrogen-phosphate

Dissolve 10 g (0.032 mole) of 1-[4-(4-chlorophenyl)-butyl]-3-propylimidazolium chloride in water and pour onto column containing 50 g anion exchange resin (BIO-RAD hydroxide form AG-15 (1-X8). Elute column with water and collect eluates having pH>9. Wash these eluates with 50 ml methylene chloride. Titrate sample to pH=5 with 10% phosphoric acid and from pH=5 to pH=4.5 with 1% phosphoric acid. Evaporate aqueous solution to dryness and 20 recrystallize from isopropyl alcohol to provide the title compound.

NMR (D_2O): δ =1.04(t,3), 1.35-2.38(m,6), 2.70(t,2), 4.12-4.50(m,4), 7.21(s,4), 7.43-7.66(m,2) and 7.80(br s,1)ppm.

25 EXAMPLE 19

A. 1-Cyclohexylmethyl-3-[4-(4-chlorophenyl)butyl]imidazolium chloride

Add 6.55 g (0.03 mole) of 1-chloro-4-(4-chlorobutyl)-benzene to 4.81 g (0.293 mole) of 1-cyclohexylmethyl-1H-imidazole and heat at 130°C for 18 hours.

Follow the progress of reaction by NMR (CDCl_3). At the completion of the reaction, dissolve reaction mixture in methanol: water (1:1) mixture and then wash with 260 ml cyclohexane. Extract aqueous layer with 200 ml methylene chloride and evaporate to provide the title compound..

B. 1-Cyclohexylmethyl-3-[4-(4-chlorophenyl)butyl]imidazolium dihydrogenphosphate

Dissolve 7.58 g (0.021 mole) of 1-cyclohexylmethyl-3-[4-(4-chlorophenyl)butyl]imidazolium chloride in water and pour onto column containing 50 g anion exchange resin (BIO-RAD hydroxide form AG-1-X8). Elute sample through column with water and collect eluates having $\text{pH} \geq 9$. Wash these eluates with 50 ml methylene chloride. Titrate sample to $\text{pH}=5$ with 10% phosphoric acid and from $\text{pH}=5$ to $\text{pH}=4.5$ with 1% phosphoric acid. Evaporate aqueous solution to dryness and recrystallize from isopropyl alcohol to provide the title compound.

NMR (D_2O): $\delta=0.90-2.30(\text{m},15)$, $2.68(\text{t},2)$, $4.05-4.58(\text{m},4)$, $7.21(\text{s},4)$, $7.60(\text{br s},2)$ and $8.95(\text{br s},1)\text{ppm}$.

EXAMPLE 20

1-Heptyl-3-[4-(4-methoxyphenyl)butyl]-2,4,5-trimethylimidazolium bromide

Add 3.23 g (0.0133 mole) of 1-(4-bromobutyl)-4-methoxybenzene to 2.77 g (0.0133 mole) of 1-heptyl-2,4,5-trimethyl-1H-imidazole and heat at 125°C for two hours. Follow the progress of the reaction by thin-layer chromatography on silica gel (methanol: 1M sodium chloride, 95:5). At the completion of the reaction, dissolve the cooled reaction mixture in 100 ml water and then wash with 100 ml cyclohexane. Extract aqueous layer with 150 ml methylene chloride and evaporate to provide the title compound.

NMR (CDCl_3): $\delta=0.69-2.07(\text{m},17)$, $2.30(2,6)$, $2.60(\text{t},2)$, $2.82(2,3)$, $3.80(\text{s},3)$, $3.95-4.35(\text{m},4)$ and $6.70-7.20(\text{m},4)\text{ppm}$.

EXAMPLE 211-[4-(4-Chlorophenyl)butyl]-3-methylimidazolium chloride

Combine 6.1 g (0.03 mole) 1-chloro-4-(4-chlorobutyl)-benzene with 2.5 g (0.03 mole) 1-methyl-1H-imidazole and heat at 120°C, following the course of the reaction by thin-layer chromatography on silica gel (methanol: 1M sodium chloride, 95:5). At the completion of the reaction, cool and triturate with ether, collect the solid and recrystallize from 50 ml hot acetone to obtain the title compound.

NMR (CDCl₃): δ=1.35-2.30(m,4), 2.60(t,2), 4.05(s,3), 4.35(t,2), 6.80-7.80(m,6) and 10.55 (s,1)ppm.

EXAMPLE 221-[4-(2,3-Dichlorophenyl)-2-butenyl]-3-heptylimidazolium 4-methylbenzenesulfonate

5.0 g (0.021 mole) 1-chloro-4-(2,3-dichlorophenyl)-2-butene and 3.5 g (0.021 mole) 1-heptyl-1H-imidazole are dissolved in 40 ml acetonitrile and heated at reflux for about 24 hours. The reaction is followed by TLC on silica gel (EM) in acetonitrile: ammonium hydroxide, 9:1, visualized by UV and iodoplatinate. The solvent is evaporated and the residue is taken up in 100 ml H₂O and washed with 4 x 35 ml ether. To the aqueous layer is added 15 ml saturated aqueous NaCl, and the aqueous layer is extracted with 3 x 100 ml methylene chloride. The combined organic layers are dried over magnesium sulfate, filtered and stripped to yield an oil. The oil is chromatographed on a column prepared from a ten-fold excess of anion exchange resin (BIO-RAD 1-X8, 20-50 mesh, hydroxide form) and eluted with deionized water. The basic fractions are pooled and titrated to pH 7 with a concentrated aqueous solution of p-toluenesulfonic acid. The water is evaporated by azeotroping with acetonitrile and the resulting oil is crystallized from acetone-ether to yield the title compound.

NMR (CDCl₃): δ =0.54-1.45 (m, 11), 1.45-2.0 (m, 2), 2.31 (s, 3), 3.50 (d, 2), 4.15 (t, 2), 4.83 (d, 2), 5.52-6.27 (m, 2), 6.95-7.53 (m, 7), 7.75 (br s, 1), 7.87 (br s, 1) and 9.80 (br s, 1) ppm.

EXAMPLE 23

(±)-1-[2-Hydroxy-2-[4-[(methanesulfonyl)amino]phenyl]ethyl]-3-methylimidazolium iodide

Heat a solution of 4.0 g (14.2 mmole) of (±)-N-[4-[1-hydroxy-2-(1H-imidazol-1-yl)ethyl]phenyl]methanesulfonamide, 2 ml (32.1 mmole) of iodomethane and 50 ml methanol in a glass pressure tube to 80°C for 24 hours. Remove the solvent *in vacuo*. Crystallize the residue from acetone. Recrystallize from acetone to provide the title compound.

NMR (CF₃CO₂D): δ =3.2 (s, 3), 4.0 (s, 3), 4.5-5.0 (m, 2), 5.3-5.6 (m, 1), 7.2-7.6 (m, 6) and 8.8 (m, 1) ppm.

EXAMPLE 24

In a manner similar to Example 16 using acetonitrile the following compounds are prepared:

- (a) 4-chloro-1-(4-chlorophenyl)-(E)-1-butene + 1-heptyl-1H-imidazole=1-[4-(4-chlorophenyl)-(E)-3-butenyl]-3-heptylimidazolium chloride.
- (b) 4-chloro-1-methylbutanol 4-methylbenzenesulfonate (Ester) + 1-heptyl-1H-imidazole=1-[4-(4-chlorophenyl)-1-methylbutyl]-3-heptylimidazolium 4-methylbenzenesulfonate.

EXAMPLE 25

In a manner similar to Example 18 the following compounds are prepared:

- (a) 1-(5-bromopentyl)-4-chlorobenzene + 1-heptyl-1H-imidazole=1-[5-(4-chlorophenyl)pentyl]-3-heptylimidazolium dihydrogen-phosphate.

- 0131302
- (b) 1-chloro-4-(2-chlorophenyl)-2-butene + 1-heptyl-1*H*-imidazole = 1-[4-(2-chlorophenyl)-2-butenyl]-3-heptylimidazolium dihydrogenphosphate.
- (c) 1-chloro-4-(4-chlorobutyl)benzene + 1-butyl-1*H*-imidazole = 1-butyl-3-[4-(4-chlorophenyl)butyl]imidazolium dihydrogenphosphate.
- (d) 1-chloro-4-(4-chlorobutyl)benzene + 1-[4-(4-chlorophenyl)-butyl-1*H*-imidazole = 1,3-bis[4-(4-chlorophenyl)butyl]-imidazolium dihydrogenphosphate.
- (e) 4-(4-chlorophenyl)-3-butyne-1-ol 4-methylbenzenesulfonate (Ester) + 1-heptyl-1*H*-imidazole = 1-[4-(4-chlorophenyl)-3-butyne-1-yl]-3-heptylimidazolium 4-methylbenzenesulfonate.

EXAMPLE 26

In a manner similar to Example 20 with further conversion to the dihydrogenphosphate the following compound is prepared:

- (a) 4-[(methanesulfonyl)amino]benzenebutanol methanesulfonate (Ester) + 1-heptyl-1*H*-imidazole = 1-heptyl-3-[4-[4-[(methylsulfonyl)amino]phenyl]butyl]imidazolium dihydrogenphosphate.

EXAMPLE 27

(E)-1-[4-(4-Fluorophenyl)-3-butenyl]-3-heptylimidazolium dihydrogenphosphate

Combine 4.62 g (0.025 mole) of 4-(4-fluorophenyl)-3-butenyl chloride and 4.16 g (0.025 mole) of 1-heptyl-1*H*-imidazole and heat at 120°C for about 8 hours, following the reaction by thin-layer chromatography. Dissolve the reaction in water and extract three times with ether. Then extract the aqueous portions with methylene chloride (10 x 30 ml). Wash the methylene chloride with water, dry over magnesium sulfate and then evaporate off the organic solvent. The residue is converted to the title compound as exemplified in Example 18 B.

NMR(D₂O): δ =0.86(t,3), 0.92-1.14(m,6), 1.14-1.30(m,2), 1.60-1.75(m,2), 2.80(q,2), 4.18(t,2), 4.46(t,2), 5.14-6.34(m,2), 7.11(dd,2), 7.40(dd,2), 7.52(s,1), 7.62(s,1) and 8.84(s,0.7)ppm.

EXAMPLE 28

(Z)-1-[4-(4-Chlorophenyl)-2-butenyl]-3-heptylimidazolium 4-methylbenzenesulfonate

5 Add 3.37 g (0.01 mole) of (Z)-4-(4-chlorophenyl)-2-buten-1-ol 4-methylbenzenesulfonate (ester) and 1.50 g (0.009 mole) of 1-heptyl-1*H*-imidazole in 40 ml of acetonitrile and stir at room temperature for two days. Evaporate the acetonitrile and triturate the residue with hexanes and ethyl acetate to yield a solid. Recrystallize from ethyl acetate to provide the
10 title compound.

EXAMPLE 29

(Z)-1-[4-(4-Chlorophenyl)-2-butenyl]-3-heptylimidazolium dihydro-genphosphate

15 Dissolve 2.7 g (0.005 mole) of (Z)-1-[4-(4-chlorophenyl)-2-butenyl]-3-heptylimidazolium 4-methylbenzenesulfonate in 25 ml of methanol/water (80/20) mixture and pour onto column containing 8 g of anion exchange resin (BIO-RAD AG-1-X8, hydroxide form). Elute the column with water, and collect eluate having pH>8. Wash the eluate with ether. Titrate the aqueous solution to
20 pH=5.0 with 10% phosphoric acid and from pH=5 to pH=4.5 with 1% phosphoric acid. Lyophilize to dryness and recrystallize from acetonitrile to provide the title compound.

NMR(D₂O): δ =0.94(t,3), 1.32(br s,8), 1.80-1.97(m,2),
3.63(d,2), 4.26(t,2), 5.05(d,2), 5.95-
25 6.08(m,1), 6.19-6.34(m,1), 7.27(d,2),
7.40(d,2), 7.60+7.61(2s,2) and 8.76
(s,1)ppm.

EXAMPLE 303-Methyl-1-[2-[4-((methylsulfonyl)amino)phenyl]-2-oxoethyl]-1H-imidazolium chloride

Heat a mixture of 160 g (0.646 mole) of N-[4-(2-chloro-1-oxoethyl)phenyl]methanesulfonamide, 55.7 g (0.678 mole) of 1-methyl-1H-imidazole and 2.1 L of acetonitrile at reflux for about 16 hours. Cool the mixture to room temperature and collect the solid and wash with 2 L of acetonitrile. Drying provides the title compound, which may be recrystallized from aqueous ethanol.

NMR (DMSO-d₆): δ = 3.14(s,3), 3.95(s,3),
5.98(s,2), 7.36(d,2),
7.69(t,2), 7.76(t,1),
8.02(d,2), 9.06(m,1), and
10.66(br s,1)ppm.

EXAMPLE 31

In a manner similar to EXAMPLE 30 the following compounds may be prepared:

(a) N-[4-(2-chloro-1-oxoethyl)phenyl]methanesulfonamide + 1,2-dimethyl-1H-imidazole = 2,3-dimethyl-1-[2-[4-((methylsulfonyl)amino)phenyl]-2-oxoethyl]-1H-imidazolium chloride

(b) N-[4-(2-chloro-1-oxoethyl)phenyl]methane-sulfonamide + 1-ethyl-1H-imidazole = 3-ethyl-1-[2-[4-((methylsulfonyl)amino)phenyl]-2-oxoethyl]-1H-imidazolium chloride

5 (c) N-[4-(2-chloro-1-oxoethyl)phenyl]methane-sulfonamide + 1-(1-methylethyl-1H-imidazole = 3-(1-methylethyl)-1-[2-[4-((methylsulfonyl)amino)phenyl]-2-oxoethyl]-1H-imidazolium chloride

10 (d) N-[4-(2-chloro-1-oxoethyl)phenyl]ethane-sulfonamide + 1-methyl-1H-imidazole = 1-[2-[4-((ethylsulfonyl)amino)phenyl]-2-oxoethyl]-3-methyl-1H-imidazolium chloride

15 (e) N-[4-(3-chloro-1-oxopropyl)phenyl]methane-sulfonamide + 1-methyl-1H-imidazole = 3-methyl-1-[3-[4-((methylsulfonyl)amino)phenyl]-3-oxopropyl]-1H-imidazolium chloride

EXAMPLE 32

(±)-1-[2-Hydroxy-2-[4-((methylsulfonyl)amino)phenyl]-ethyl]-3-methyl-1H-imidazolium chloride

20 A solution of 152.3 g (0.463 mole) of 3-methyl-1-[2-[4-((methylsulfonyl)amino)phenyl]-2-oxoethyl]-1H-imidazolium chloride in 1 L of distilled water and 7.62 g 10% Pd/C catalyst are placed in a 2 L Parr bottle and hydrogenated at 50 psi pressure for ca. 5 hours.

25 Additional H₂ is added as needed to maintain 50 psi

5 NMR (DMSO-d₆): δ = 2.98(s,3), 3.87(s,3),
4.11(m,1), 4.22(m,1),
4.91(m,1), 6.10(d,1),
7.23(d,2), 7.37(d,2),
7.71(d,2), 9.16(s,1), and
10 9.84(s,1)ppm.

In a manner similar to EXAMPLE 32 the following compounds are prepared:

15 (a) 2,3-dimethyl-1-[2-[4-((methylsulfonyl)amino)-phenyl]-2-oxoethyl]-1H-imidazolium chloride + H₂ = (±)-2,3-dimethyl-1-[2-hydroxy-2-[4-((methylsulfonyl)amino)-phenyl]ethyl]-1H-imidazolium chloride

(b) 3-ethyl-1-[2-[4-((methylsulfonyl)amino)phenyl]-2-oxoethyl]-1H-imidazolium chloride + H₂ = (±)-3-ethyl-1-[2-hydroxy-2-[4-((methylsulfonyl)amino)phenyl]ethyl]-1H-imidazolium chloride

(c) 3-(1-methylethyl)-1-[2-[4-((methylsulfonyl)-amino)phenyl]-2-oxoethyl]-1H-imidazolium chloride + H₂ =
([±])-1-[2-hydroxy-2-[4-((methylsulfonyl)amino)phenyl]-ethyl]-3-(1-methylethyl)-1H-imidazolium chloride

(d) 1-[2-[4-((ethylsulfonyl)amino)phenyl]-2-oxo-ethyl]-3-methyl-1H-imidazolium chloride + H₂ = (±)-1-[2-[4-((ethylsulfonyl)amino)phenyl]-2-hydroxyethyl]-3-methyl-1H-imidazolium chloride

5 (e) 3-methyl-1-[3-[4-((methylsulfonyl)amino)-phenyl]-3-oxopropyl]-1H-imidazolium chloride + H₂ = (±)-1-[3-hydroxy-3-[4-((methylsulfonyl)amino)phenyl]propyl]-3-methyl-1H-imidazolium chloride

EXAMPLE 34

10 1-[2-(4-Aminophenyl)ethyl]-1H-imidazole

A solution of 10 g (0.046 mole) of 1-[2-(4-nitrophenyl)ethyl]-1H-imidazole in 100 mL of ethanol, 50 mL of 10% aqueous hydrochloric acid and 1.0 g of 5% Pd/C catalyst are placed in a small Parr bottle and
15 hydrogenated at 50 psi pressure for about 24 hours. Additional H₂ is added as needed to maintain 50 psi pressure. The reaction mixture is filtered and the solvent is concentrated in vacuo. The residue is
20 dissolved in 100 mL of water and made basic (pH 9) with potassium carbonate. Extract the aqueous mixture with methylene chloride (2x100 mL) and combine. Evaporation of the solvent yields the title compound.

EXAMPLE 351-[2-[4-[(Methylsulfonyl)amino]phenyl]ethyl]-1H-imidazole

5 To a -10°C solution of 8 g (0.064 mole) of 1-[2-(4-aminophenyl)ethyl]-1H-imidazole and 10 mL triethylamine in 200 mL of methylene chloride add 5 mL of methanesulfonyl chloride dropwise. Stir the resulting solution at -10°C for about 30 minutes then warm to ambient temperature for about 30 minutes. Wash the reaction mixture with water (2x100 mL). Extract with 10% aqueous sodium hydroxide (2x50 mL). The combined aqueous extracts are neutralized with ammonium chloride then extracted with methylene chloride (2x100 mL) and combined. Remove the solvent in vacuo to provide the title compound, which may be recrystallized from tetrahydrofuran.

15 NMR (CF₃COOH): δ = 3.1-3.5(m,5), 4.4-4.8(t,2), 7.1-7.6(m,6), and 8.4(s,1)ppm.

EXAMPLE 363-Methyl-1-[2-[4-[(methylsulfonyl)amino]phenyl]ethyl]-1H-imidazolium iodide

20 Heat a mixture of 5 g (0.019 mole) of N-[4-[2-(1H-imidazol-1-yl)ethyl]phenyl]methanesulfonamide, 5 mL of iodomethane and 25 mL of methanol in a pressure tube at 75°C for about 24 hours. Cool the mixture to room

temperature and remove the solvents in vacuo. Slurry the residue in acetone (50 mL) and filter and wash with acetone (100 mL). Drying provides the title compound, which may be recrystallized from methanol.

5 NMR (CF_3COOD): δ = 3.15-3.55(m,5), 4.05(s,3),
4.45-4.85(t,2), 7.20-7.55(m,6),
and 8.65(s,1)ppm.

EXAMPLE 37

10 (\pm)-1-[2-Hydroxy-2-[4-((methylsulfonyl)amino)phenyl]ethyl]-1H-imidazole

Heat a mixture of 30.83 g (0.097 mole) of (\pm)-1-[2-hydroxy-2-[4-((methylsulfonyl)amino)phenyl]ethyl]-1H-imidazole hydrochloride and 5.24 g (0.097 mole) of sodium methylate in 100 mL of methanol and filter hot through 300 g of alumina (Fisher, neutral, activity III). Wash the alumina with 20% methanol in methylene chloride (200 mL). Concentrate the filtrate in vacuo to afford the title compound.

20 NMR ($\text{DMSO}-d_6$): δ = 2.95(s,3), 4.03(m,1),
4.13(m,1), 4.79(m,1),
5.69(d,1), 6.82(s,1),
7.10(s,1), 7.16(d,2),
7.28(d,2), 7.47(s,1), and
9.74(s,1)ppm.
25 7.70(s,1), 9.02(s,1),
9.77(s,1) and 14.30(br s,1)ppm.

$[\alpha]_D^{23} = +14.9^\circ$ (C, 1.95, 2N aqueous NaOH)

EXAMPLE 38

(+)-1-[2-Hydroxy-2-[4-((methylsulfonyl)amino)phenyl]ethyl]-1H-imidazole-d-7,7-dimethyl-2-oxobicyclo[2.2.1]heptane-1-methanesulfonate

5 Heat a mixture of 27.3 g (0.097 mole) of (+)-1-[2-hydroxy-2-[4-((methylsulfonyl)amino)phenyl]ethyl]-1H-imidazole, 22.5 g (0.097 mole) of d-7,7-dimethyl-2-oxobicyclo[2.2.1]heptane-1-methanesulfonic acid in 200 mL of water and 200 mL of ethanol to effect complete
10 solution, then remove the solvent in vacuo. Recrystallize the residue from methanol and recrystallize the resulting crystals 3 more times from aqueous methanol to provide the title compound.

15 NMR (DMSO-d₆): δ = 0.75(s,3), 1.05(s,3),
1.26-1.29(m,2), 1.83-1.93(m,2),
1.97(t,1), 2.20(m,1),
2.37(d,1), 2.98(m,1),
2.87(d,1), 2.98(s,3),
4.20-4.45(m,2), 4.92(br d,1),
20 5.92(br s,1), 7.20(d,2),
7.35(d,2), 7.65(s,1),
7.70(s,1), 9.02(s,1),
9.77(s,1) and 14.30(br s,1)ppm.

$[\alpha]_D^{23} = +14.9^\circ$ (C, 1.95, 2N aqueous NaOH)

EXAMPLE 39

(-)-1-[2-Hydroxy-2-[4-((methylsulfonyl)amino)phenyl]ethyl]-1H-imidazole-1-7,7-dimethyl-2-oxobicyclo[2.2.1]heptane-1-methanesulfonate

Heat a mixture of 11.05 g (0.039 mole) of (+)-1-[2-hydroxy-2-[4-((methylsulfonyl)amino)phenyl]ethyl]-1H-imidazole, 9.12 g (0.039 mole) of 1-7,7-dimethyl-2-oxobicyclo[2.2.1]heptane-1-methanesulfonic acid in 100 mL of water and 100 mL of ethanol to effect complete solution, then remove the solvent in vacuo. Recrystallize the residue from methanol and recrystallize the resulting crystals 3 more times from aqueous methanol to provide the title compound.

NMR (DMSO- d_6): δ = 0.74(s,3), 1.05(s,3),
 1.24-1.29(m,2), 1.76-1.82(m,2),
 1.93(t,1), 2.20(m,1),
 2.37(d,1), 2.67(m,1),
 2.87(d,1), 2.98(s,3),
 4.22-4.40(m,2), 4.92(br d,1),
 5.96(br s,1), 7.20(d,2),
 7.34(d,2), 7.65(s,1),
 7.69(s,1), 9.01(s,1),
 9.77(s,1), and
 14.30(br s,1)ppm.

$[\alpha]_D^{23}$ = -14.9° (C, 1.95, 2N aqueous NaOH).

EXAMPLE 40

In a manner similar to Example 37, the following compounds may be prepared:

5 (a) (+)-1-[2-hydroxy-2-[4-((methylsulfonyl)amino)-phenyl]ethyl]-1H-imidazole-2,7,7-dimethyl-2-oxobicyclo-[2.2.1]heptane-1-methanesulfonate + sodium methylate = (-)-1-[2-hydroxy-2-[4-((methylsulfonyl)amino)phenyl]-ethyl]-1H-imidazole

$[\alpha]_D^{23} = -59.5^\circ$ (C, 1.715, 1N aqueous HCl).


10 (b) (-)-1-[2-hydroxy-2-[4-((methylsulfonyl)amino)-phenyl]ethyl]-1H-imidazole-2,7,7-dimethyl-2-oxobicyclo-[2.2.1]heptane-1-methanesulfonate + sodium methoxide = (+)-1-[2-hydroxy-2-[4-((methylsulfonyl)amino)phenyl]-ethyl]-1H-imidazole


15 $[\alpha]_D^{23} = +59.7^\circ$ (C, 1.740, 1N NaOH)


We Claim:

-1-

A compound of the formula:



10 wherein R = straight or branched chain alkyl having 1-12 carbon atoms, cycloalkyl(lower)alkyl, $-(CH_2)_t-$ , or



R_1, R_2, R_3 = hydrogen, a straight chain alkyl group of 1-4 carbon atoms,

20 R_4, R_5 = hydrogen, halogen, hydroxy, lower alkyl, lower alkoxy, cyano, nitro, carbamoyl, lower alkylcarbamoyl, lower alkanoylamido, sulfamoyl, substituted sulfamoyl, sulfamoylamino, N-lower alkylsulfamoylamino, lower alkylsulfonamido, substituted lower alkylsulfonamido, trifluoromethanesulfonamido, ureido, N-lower alkylureido
25 or when taken together on adjacent carbon atoms of the ring to which they are attached may be joined to form the methylene dioxy moiety;

30 R_6 = hydrogen, halogen, hydroxy, straight chain alkyl of 1-6 carbon atoms, lower alkoxy, cyano and carbamoyl;

0131302

A = phenyl, furanyl, pyridinyl, pyrrolyl, thiazolyl, oxazolyl, the benzo derivatives of the foregoing and thiadiazolyl;

n = 0-2,

p = 0-5,

t = 1-5,

Y = methylene, hydroxymethylene, arylmethylene, substituted arylmethylene, phenylene, ethynylene, $\overset{R_7 R_8}{-C=C-}$,



$-SO_2NH-$ wherein R_7 , R_8 is hydrogen, C_1-C_4 lower alkyl or taken together form a 5 or 6 membered ring;


R_9 , R_{10} is C_1-C_4 lower alkyl, or taken together may form a 5 or 6 membered ring or either may by

hydrogen if the other is C_1-C_4 alkyl; R_{11} is hydrogen, lower alkyl, cyano and carbamoyl,

with the provisos that:

(a) when Y is $>N-R_{11}$, $-SO_2NH-$, $-NHSO_2-$, then p must be an integer of 2-5,

(b) when Y is hydroxymethylene then R cannot be



(c) when Y is methylene, ethynylene, $\overset{R_7 R_8}{-C=C-}$, $=C=C$

then p must be an integer of 1-5,

(d) when the sum of the carbon atoms in the chain n, Y, p is 2 then both R_4 and R_5 cannot be hydrogen,

(e) when Y is carbonyl then both R_4 and R_5 cannot be hydrogen;

Z^- = halide, tosylate, sulfate, phosphate, methane-sulfonate, and the molecular compounds of the foregoing quaternary salts with aromatic dicarboxylic acids.

-2-

A compound according to claim 1 wherein said aromatic dicarboxylic acids are selected from the group consisting of phthalic acid, 1,8-naphthalenedicarboxylic acid, 1,2-naphthalenedicarboxylic acid and 2,3-naphthalenedicarboxylic acid.

-3-

A compound according to claim 1 wherein A is phenyl and one of R₁ or R₂ is other than hydrogen.

-4-

A compound according to claim 3 wherein R is C₇H₁₅.

-5-

A compound according to claim 3 wherein R is CH₃ and/or R₁ is CH₃.

-6-

A compound of claim 4 which is 1-[4-(4-chlorophenyl)butyl]-3-heptylimidazolium chloride (1:1) compound with phthalic acid.

-7-

A compound of claim 4 which is 1-heptyl-3-[4-(4-methoxyphenyl)butyl]imidazolium bromide.

-8-

A compound of claim 1 which is 1-heptyl-3-(4-phenylbutyl)imidazolium chloride 1:1 complex with phthalic acid.

-9-

A compound of claim 4 which is 1-[4-[4-(acetylamino)phenyl]butyl]-3-heptylimidazolium 4-methylbenzenesulfonate.

-10-

A compound of claim 1 which is 1-[4-(4-chlorophenyl)butyl]-3-propylimidazolium dihydrogenphosphate.

-11-

A compound of claim 4 which is 1-[4-(4-fluorophenyl)butyl]-3-heptylimidazolium dihydrogenphosphate.

-12-

5 A compound of claim 1 which is 1-[4-(4-chlorophenyl)butyl]-3-pentylimidazolium dihydrogenphosphate.

-13-

A compound of claim 1 which is 1-butyl-3-[4-(4-chlorophenyl)-butyl]imidazolium dihydrogenphosphate.

10

-14-

A compound of claim 4 which is 1-heptyl-3-[4-[4-[(methylsulfonyl)amino]phenyl]butyl]imidazolium dihydrogenphosphate .

-15-

15 A compound of claim 4 which is 1-[4-(4-chlorophenyl)-1-methylbutyl]-3-heptylimidazolium 4-methylbenzenesulfonate.

-16-

A compound of claim 4 which is 1-[4-(4-chlorophenyl)-3-butyryl]-3-heptylimidazolium 4-methylbenzenesulfonate.

-17-

20 A compound of claim 5 which is 1,2-Dimethyl-3-[4-[4-[(methylsulfonyl)amino]phenyl]butyl]imidazolium methanesulfonate.

-18-

A compound of claim 5 which is 1-[4-(4-chlorophenyl)butyl]-2,3-dimethylimidazolium chloride.

25

-19-

A compound of claim 1 which is 1-[2-hydroxy-2-[4-[(methanesulfonyl)amino]phenyl]ethyl]-3-methylimidazolium iodide.

-20-

30 A compound of claim 4 which is 1-heptyl-3-[2-[4-[(methanesulfonyl)amino]phenyl]-2-oxoethyl]imidazolium bromide.

-21-

A compound of claim 4 which is (Z)-1-[4-(4-chlorophenyl)-2-butenyl]-3-heptylimidazolium dihydrogenphosphate.

-22-

5 A compound of claim 1 which is


1-Methyl-3-[2-[4-((methylsulfonyl)-amino)phenyl]-ethyl]-1H-imidazolium iodide.

-23-

10 1-Methyl-3-(2-(4-((methylsulfonyl)amino)phenyl)-2-oxoethyl)-1H-imidazolium bromide.

-24-

15 A pharmaceutical composition for the suppression of cardiac arrhythmic in a mammalian subject which comprises administering to said subject an amount effective for the suppression of said arrhythmia of a compound of the formula



wherein T is nitro, amino, lower alkansulfonamido and Q is a straight chain alkyl of 1-4 carbon atoms, $-\overset{\text{O}}{\parallel}{\text{C}}-\text{CH}_2-$, $-\text{CHOHCH}_2-$, and the pharmaceutically acceptable acid addition salts thereof.

20

-25-

A pharmaceutical composition for the suppression of cardiac arrhythmia comprising an antiarrhythmic effective amount of a compound of claim 1 together, with a non-toxic pharmaceutically acceptable carrier.


-26-


A pharmaceutical composition according to claim 24 wherein the compound of the formula is one of:


- 1- $\left[4-(4\text{-Nitrophenyl})\text{butyl}\right]/\text{imidazole hydrochloride}$
 5 N- $\left[4-(1\text{H-Imidaz-1-ylacetyl})\text{phenyl}\right]/\text{methansulfonamide}$
 2-(1H-Imidaz-1-yl)-1-(4-nitrophenyl)ethanone hydrochloride
 N- $\left[4-(1\text{-Hydroxy-2-(1H-imidazol-1-yl)ethyl})\text{phenyl}\right]/\text{methane-sulfonamide}$
 1-(4-Aminiphenyl)-2-(1H-imidazol-1-yl)ethanone
 10 α -(4-Nitrophenyl)-1H-imidazole-1-ethanol
 N- $\left[4-(2-(1\text{H-Imidazol-1-yl)ethyl})\text{-phenyl}\right]/\text{methansulfonamide}$
 1- $\left[2-(4\text{-Nitrophenyl)ethyl}\right]/\text{1H-imidazole hydrochloride.}$

-27-

A process for preparing a compound of the formula:



wherein R = straight or branched chain alkyl having 1-12 carbon atoms, cycloalkyl (lower) alkyl, $-(\text{CH}_2)_t$ --R₆, or




- R_1, R_2, R_3 = hydrogen, a straight chain alkyl group of 1-4 carbon atoms,
- R_4, R_5 = hydrogen, halogen, hydroxy, lower alkyl, lower alkoxy, cyano, nitro, carbamoyl, lower alkylcarbamoyl, lower alkanoylamido, sulfamoyl, substituted sulfamoyl, sulfamoylamino, N-lower alkylsulfamoylamino, lower alkylsulfonamido, substituted lower alkylsulfonamido, trifluoromethanesulfonamido, ureido, N-lower alkylureido or when taken together on adjacent carbon atoms of the ring to which they are attached may be joined to form the methylene dioxy moiety:
- R_6 = hydrogen, halogen, hydroxy, straight chain alkyl of 1-6 carbon atoms, lower alkoxy, cyano and carbamoyl;
- A = phenyl, furanyl, pyridinyl, pyrrolyl, thiazolyl, oxazolyl, the benzo derivatives of the foregoing and thiadiazolyl;
- n = 0-2,
- p = 0-5,
- t = 1-5,
- Y = methylene, hydroxymethylene, arylmethylene, substituted arylmethylene, phenylene, ethynylene, $\overset{R_7 R_8}{-C=C-}$, $\overset{R_9}{\underset{|}{-CH}}-\overset{R_{10}}{\underset{|}{-CH}}-$, $\overset{R_7}{\underset{|}{C}}=\overset{R_8}{\underset{|}{C}}$, carbonyl, $>N-R_{11}$, $-NHSO_2-$, $-SO_2NH-$ wherein R_7, R_8 is hydrogen, C_1-C_4 lower alkyl or taken together form a 5 or 6 membered ring; R_9, R_{10} is C_1-C_4 lower alkyl, or taken together may form a 5 or 6 membered ring or either may be hydrogen if the other is C_1-C_4 alkyl; R_{11} is hydrogen, lower alkyl, cyano and carbamoyl,

with the provisos that:

(a) when Y is $>N-R_{11}$, $-SO_2NH-$, $-NHSO_2-$, then p must be an integer of 2-5,

(b) when Y is hydroxymethylene then R cannot be

5



(c) when Y is methylene, ethynylene, $\begin{array}{c} R_7 \quad R_8 \\ | \quad | \\ -C=C- \end{array}$, $\begin{array}{c} R_7 \\ \diagup \\ C \\ \diagdown \\ R_8 \end{array}$

then p must be an integer of 1-5,

(d) when the sum of the carbon atoms in the chain n, Y, p is 2 then both R_4 and R_3 cannot be hydrogen,

10

(e) when Y is carbonyl then both R_4 and R_3 cannot be hydrogen;


Z^- = halide, tosylate, sulfate, phosphate, methane-

sulfonate, and the molecular compounds of the foregoing quaternary salts with aromatic dicarboxylic acids,

15


comprising
reacting

(A) a corresponding compound of the formula:



with a corresponding compound of the formula

20



and reacting the resultant product with a
corresponding compound of the formula




or

5 (B) a corresponding compound of the formula




with a corresponding compound of the formula:



10

and reacting the resultant product with a
corresponding compound of the formula:



15

wherein in both reactions (A) and (B),
Q is a leaving group which is halo or tosyl
and for the resultant products of reactions (A)
and (B) which do not contain the Z group, reacting
the resultant product with an acid whose anion
is Z, and, in each case, optionally, further
conventionally reacting the resultant compound to
convert one of the groups R_4 , R_5 , Y, R_6 or Z into
another such group.

20